

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Rarely or never wore seat belts				
Female	7.8	(6.3 - 9.3)	3.8	(2.2 - 5.4) *
Male	12.5	(10.3 - 14.7)	12.5	(9.9 - 15.1)
Total	10.2	(8.4 - 12.0)	8.2	(6.7 - 9.7)
Rarely or never wore bicycle helmets				
Female	79.9	(75.9 - 83.9)	91.7	(88.0 - 95.4) *
Male	86.1	(83.3 - 88.9)	92.9	(90.3 - 95.5) *
Total	83.4	(80.2 - 86.6)	92.3	(89.9 - 94.7) *
Rode with a driver who had been drinking alcohol				
Female	29.6	(27.2 - 32.0)	21.6	(18.0 - 25.2) *
Male	27.2	(25.2 - 29.2)	27.4	(22.4 - 32.4)
Total	28.5	(26.6 - 30.4)	24.6	(21.1 - 28.1)
Drove after drinking alcohol				
Female	8.1	(7.1 - 9.1)	7.3	(4.9 - 9.7)
Male	11.7	(10.3 - 13.1)	15.0	(11.1 - 18.9)
Total	9.9	(8.9 - 10.9)	11.2	(8.5 - 13.9)
Carried a weapon				
Female	7.1	(6.3 - 7.9)	6.5	(4.6 - 8.4)
Male	29.8	(27.2 - 32.4)	31.5	(28.3 - 34.7)
Total	18.5	(16.9 - 20.1)	19.2	(16.8 - 21.6)
Carried a gun				
Female	0.9	(0.5 - 1.3)	0.9	(0.2 - 1.6)
Male	9.9	(8.5 - 11.3)	10.5	(7.9 - 13.1)
Total	5.4	(4.6 - 6.2)	5.8	(4.6 - 7.0)
In a physical fight				
Female	28.1	(26.3 - 29.9)	21.7	(17.9 - 25.5) *
Male	43.4	(41.4 - 45.4)	36.7	(33.1 - 40.3) *
Total	35.9	(34.4 - 37.4)	29.3	(26.3 - 32.3) *
Injured in a physical fight				
Female	2.4	(2.0 - 2.8)	2.4	(1.1 - 3.7)
Male	4.8	(4.1 - 5.5)	4.5	(3.3 - 5.7)
Total	3.6	(3.2 - 4.0)	3.4	(2.4 - 4.4)
Dating violence				
Female	9.3	(8.5 - 10.1)	13.5	(10.7 - 16.3) *
Male	9.0	(8.2 - 9.8)	11.6	(9.0 - 14.2)
Total	9.2	(8.6 - 9.8)	12.5	(10.5 - 14.5)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana		
	Percent	(95% CI)	Percent	(95% CI)	
Forced to have sexual intercourse					
Female	10.8	(9.6 - 12.0)	---	---	---
Male	4.2	(3.5 - 4.9)	---	---	---
Total	7.5	(6.8 - 8.2)	---	---	---
Carried a weapon on school property					
Female	2.6	(2.0 - 3.2)	3.4	(1.7 - 5.1)	
Male	10.2	(8.6 - 11.8)	8.1	(6.1 - 10.1)	
Total	6.5	(5.6 - 7.4)	5.8	(4.4 - 7.2)	
Threatened or injured w/ a weapon on school property					
Female	6.1	(5.3 - 6.9)	6.8	(4.4 - 9.2)	
Male	9.7	(8.9 - 10.5)	10.7	(7.7 - 13.7)	
Total	7.9	(7.2 - 8.6)	8.8	(6.9 - 10.7)	
Engaged in a physical fight on school property					
Female	8.8	(7.8 - 9.8)	8.0	(5.9 - 10.1)	
Male	18.2	(16.4 - 20.0)	14.4	(11.7 - 17.1)	
Total	13.6	(12.5 - 14.7)	11.2	(9.3 - 13.1)	
Felt too unsafe to go to school					
Female	6.3	(4.8 - 7.8)	4.9	(3.4 - 6.4)	
Male	5.7	(4.6 - 6.8)	3.8	(2.4 - 5.2)	
Total	6.0	(4.8 - 7.2)	4.3	(3.4 - 5.2)	
Property stolen or deliberately damaged					
Female	28.0	(25.9 - 30.1)	26.9	22.5 - 31.3	
Male	31.4	(29.8 - 33.0)	34.4	30.9 - 37.9	
Total	29.8	(28.3 - 31.3)	30.7	27.5 - 33.9	
Felt sad or hopeless					
Female	36.7	(34.8 - 38.6)	33.6	(28.4 - 38.8)	
Male	20.4	(19.1 - 21.7)	21.2	(18.1 - 24.3)	
Total	28.5	(27.3 - 29.7)	27.3	(23.9 - 30.7)	
Seriously considered attempting suicide					
Female	21.8	(20.5 - 23.1)	22.0	(18.7 - 25.3)	
Male	12.0	(11.0 - 13.0)	14.3	(11.5 - 17.1)	
Total	16.9	(16.0 - 17.8)	18.0	(15.7 - 20.3)	

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Made a suicide plan				
Female	16.2	(15.0 - 17.4)	17.0	(13.7 - 20.3)
Male	9.9	(8.7 - 11.1)	12.6	(9.7 - 15.5)
Total	13.0	(12.1 - 13.9)	14.8	(12.2 - 17.4)
Attempted suicide				
Female	10.8	(9.7 - 11.9)	11.4	(8.7 - 14.1)
Male	6.0	(4.8 - 7.2)	7.9	(5.5 - 10.3)
Total	8.4	(7.5 - 9.3)	9.6	(7.6 - 11.6)
Suicide attempt required medical attention				
Female	2.9	(2.3 - 3.5)	3.4	(2.0 - 4.8)
Male	1.8	(1.3 - 2.3)	3.6	(2.0 - 5.2)
Total	2.3	(1.9 - 2.7)	3.5	(2.4 - 4.6)
Lifetime cigarette use				
Female	52.7	(49.3 - 56.1)	54.0	(47.2 - 60.8)
Male	55.9	(53.1 - 58.7)	59.7	(55.6 - 63.8)
Total	54.3	(51.3 - 57.3)	56.9	(52.6 - 61.2)
Lifetime daily cigarette use				
Female	13.5	(11.5 - 15.5)	15.5	(11.8 - 19.2)
Male	13.3	(11.4 - 15.2)	16.8	(13.2 - 20.4)
Total	13.4	(11.5 - 15.3)	16.1	(12.9 - 19.3)
Current cigarette use				
Female	23.0	(20.4 - 25.6)	20.5	(15.8 - 25.2)
Male	22.9	(20.7 - 25.1)	23.2	(18.7 - 27.7)
Total	23.0	(20.7 - 25.3)	21.9	(17.9 - 25.9)
Current frequent cigarette use				
Female	9.3	(7.5 - 11.1)	9.7	(6.6 - 12.8)
Male	9.3	(7.8 - 10.8)	11.5	(8.1 - 14.9)
Total	9.4	(7.9 - 10.9)	10.6	(7.7 - 13.5)
Smoked > 10 cigarettes/day				
Female	7.2	(5.2 - 9.2)	10.6	(6.0 - 15.2)
Male	14.2	(12.1 - 16.3)	15.3	(9.0 - 21.6)
Total	10.7	(9.0 - 12.4)	13.2	(9.2 - 17.2)
Tried to quit smoking cigarettes				
Female	60.3	(56.5 - 64.1)	67.5	(60.4 - 74.6)
Male	48.9	(45.8 - 52.0)	55.7	(48.6 - 62.8)
Total	54.6	(52.2 - 57.0)	61.1	(56.0 - 66.2)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Purchased cigarettes at a store or gas station				
Female	11.7	(8.3 - 15.1)	13.3	(4.1 - 22.5)
Male	18.8	(15.5 - 22.1)	22.8	(16.7 - 28.9)
Total	15.2	(12.5 - 17.9)	18.2	(12.5 - 23.9)
Current smokeless tobacco use				
Female	2.2	(1.6 - 2.8)	2.1	(1.0 - 3.2)
Male	13.6	(11.1 - 16.1)	14.8	(11.1 - 18.5)
Total	8.0	(6.6 - 9.4)	8.6	(6.6 - 10.6)
Current cigar use				
Female	8.7	(7.5 - 9.9)	8.3	(5.8 - 10.8)
Male	19.2	(17.2 - 21.2)	22.7	(18.9 - 26.5)
Total	14.0	(12.5 - 15.5)	15.6	(12.8 - 18.4)
Current tobacco use				
Female	25.1	(22.5 - 27.7)	22.8	(18.0 - 27.6)
Male	31.7	(28.7 - 34.7)	35.4	(30.7 - 40.1)
Total	28.4	(25.7 - 31.1)	29.2	(25.2 - 33.2)
Lifetime alcohol use				
Female	74.8	(70.9 - 78.7)	73.9	(69.7 - 78.1)
Male	73.8	(71.1 - 76.5)	76.1	(72.7 - 79.5)
Total	74.3	(71.2 - 77.4)	75.0	(71.8 - 78.2)
Current alcohol use				
Female	42.8	(39.7 - 45.9)	41.3	(36.0 - 46.6)
Male	43.8	(41.1 - 46.5)	41.6	(37.0 - 46.2)
Total	43.3	(40.6 - 46.0)	41.4	(37.2 - 45.6)
Episodic heavy drinking				
Female	23.5	(21.2 - 25.8)	21.7	(16.6 - 26.8)
Male	27.5	(24.9 - 30.1)	27.4	(23.1 - 31.7)
Total	25.5	(23.3 - 27.7)	24.6	(20.5 - 28.7)
Lifetime marijuana use				
Female	35.9	(33.0 - 38.8)	35.1	(29.9 - 40.3)
Male	40.9	(38.3 - 43.5)	41.3	(36.9 - 45.7)
Total	38.4	(35.9 - 40.9)	38.2	(34.4 - 42.0)
Current marijuana use				
Female	18.2	(16.3 - 20.1)	16.7	(13.7 - 19.7)
Male	22.1	(20.2 - 24.0)	21.0	(17.6 - 24.4)
Total	20.2	(18.6 - 21.8)	18.9	(16.2 - 21.6)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Lifetime cocaine use				
Female	6.8	(5.8 - 7.8)	5.8	(3.4 - 8.2)
Male	8.4	(7.1 - 9.7)	7.8	(5.4 - 10.2)
Total	7.6	(6.6 - 8.6)	6.8	(4.7 - 8.9)
Current cocaine use				
Female	2.8	(2.1 - 3.5)	2.3	(1.1 - 3.5)
Male	4.0	(3.1 - 4.9)	3.6	(2.0 - 5.2)
Total	3.4	(2.8 - 4.0)	3.0	(1.9 - 4.1)
Lifetime injecting illegal drug use				
Female	1.1	(0.7 - 1.5)	1.6	(0.4 - 2.8)
Male	3.0	(2.5 - 3.5)	2.5	(1.3 - 3.7)
Total	2.1	(1.8 - 2.4)	2.1	(1.2 - 3.0)
Lifetime inhalant use				
Female	13.5	(12.0 - 15.0)	12.7	(9.1 - 16.3)
Male	11.3	(10.0 - 12.6)	15.4	(12.5 - 18.3)
Total	12.4	(11.1 - 13.7)	14.1	(11.5 - 16.7)
Lifetime illegal steroid use				
Female	3.2	(2.7 - 3.7)	4.3	(2.3 - 6.3)
Male	4.8	(4.0 - 5.6)	5.3	(3.5 - 7.1)
Total	4.0	(3.5 - 4.5)	4.8	(3.2 - 6.4)
Lifetime heroine use				
Female	1.4	(1.0 - 1.8)	1.5	(0.3 - 2.7)
Male	3.3	(2.7 - 3.9)	3.0	(1.6 - 4.4)
Total	2.4	(2.0 - 2.8)	2.3	(1.2 - 3.4)
Lifetime methamphetamine use				
Female	6.0	(4.8 - 7.2)	6.1	(3.9 - 8.3)
Male	6.3	(5.3 - 7.3)	7.9	(5.2 - 10.6)
Total	6.2	(5.3 - 7.1)	7.0	(5.2 - 8.8)
Lifetime ecstasy use				
Female	5.3	(4.5 - 6.1)	4.9	(2.7 - 7.1)
Male	7.2	(6.0 - 8.4)	7.9	(4.7 - 11.1)
Total	6.3	(5.4 - 7.2)	6.4	(4.0 - 8.8)
Smoked a whole cigarette before age 13 years				
Female	13.6	(12.1 - 15.1)	15.4	(12.3 - 18.5)
Male	18.3	(16.5 - 20.1)	18.2	(14.8 - 21.6)
Total	16.0	(14.5 - 17.5)	16.8	(13.9 - 19.7)
Drank alcohol before age 13 years				
Female	22.0	(20.1 - 23.9)	17.6	(14.0 - 21.2)
Male	29.2	(26.7 - 31.7)	25.9	(22.0 - 29.8)
Total	25.6	(23.9 - 27.3)	21.8	(18.7 - 24.9)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Tried marijuana before age 13 years				
Female	6.3	(5.5 - 7.1)	6.5	(3.5 - 9.5)
Male	11.0	(9.6 - 12.4)	10.6	(7.7 - 13.5)
Total	8.7	(7.8 - 9.6)	8.6	(6.3 - 10.9)
Cigarette use on school property				
Female	6.2	(5.0 - 7.4)	5.0	(2.8 - 7.2)
Male	7.4	(6.6 - 8.2)	7.2	(4.9 - 9.5)
Total	6.8	(6.0 - 7.6)	6.1	(4.3 - 7.9)
Smokeless tobacco use on school property				
Female	0.8	(0.5 - 1.1)	0.6	(-0.1 - 1.3)
Male	9.2	(7.0 - 11.4)	7.8	(5.9 - 9.7)
Total	5.0	(3.8 - 6.2)	4.3	(3.2 - 5.4)
Alcohol use on school property				
Female	3.3	(2.7 - 3.9)	1.9	(0.6 - 3.2)
Male	5.3	(4.5 - 6.1)	4.8	(2.8 - 6.8)
Total	4.3	(3.7 - 4.9)	3.4	(2.2 - 4.6)
Marijuana use on school property				
Female	3.0	(2.4 - 3.6)	1.2	(0.6 - 1.8) *
Male	6.0	(5.1 - 6.9)	5.6	(3.6 - 7.6)
Total	4.5	(3.9 - 5.1)	3.4	(2.3 - 4.5)
Offered, sold or given illegal drug on school property				
Female	21.8	(19.8 - 23.8)	24.7	(22.4 - 27.0)
Male	28.8	(26.4 - 31.2)	33.0	(29.4 - 36.6)
Total	25.4	(23.3 - 27.5)	28.9	(26.3 - 31.5)
Ever had sexual intercourse				
Female	45.7	(42.1 - 49.3)	43.0	(37.7 - 48.3)
Male	47.9	(44.5 - 51.3)	46.0	(41.2 - 50.8)
Total	46.8	(43.5 - 50.1)	44.5	(40.7 - 48.3)
Had first sexual intercourse before age 13				
Female	3.7	(3.0 - 4.4)	---	---
Male	8.8	(7.7 - 9.9)	---	---
Total	6.2	(5.4 - 7.0)	---	---
Had > or = to 4 sex partners during lifetime				
Female	12.0	(10.4 - 13.6)	---	---
Male	16.5	(14.7 - 18.3)	---	---
Total	14.3	(12.8 - 15.8)	---	---

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Currently sexually active				
Female	34.6	(31.6 - 37.6)	34.2	(29.8 - 38.6)
Male	33.3	(30.7 - 35.9)	35.0	(31.3 - 38.7)
Total	33.9	(31.4 - 36.4)	34.6	(31.4 - 37.8)
Condom use during last sexual intercourse				
Female	55.9	(53.1 - 58.7)	62.6	(56.0 - 69.2)
Male	70.0	(66.9 - 73.1)	62.6	(56.0 - 69.2)
Total	62.8	(60.7 - 64.9)	62.6	(57.2 - 68.0)
Birth control pill use before last sexual intercourse				
Female	20.6	(16.9 - 24.3)	---	---
Male	14.6	(12.1 - 17.1)	---	---
Total	17.6	(15.0 - 20.2)	---	---
Alcohol or drug use before last sexual intercourse				
Female	19.0	(17.0 - 21.0)	---	---
Male	27.6	(24.4 - 30.8)	---	---
Total	23.3	(21.1 - 25.5)	---	---
Taught about AIDS or HIV infection in school				
Female	88.5	(86.6 - 90.4)	93.5	(91.8 - 95.2) *
Male	87.2	(85.2 - 89.2)	90.4	(87.9 - 92.9)
Total	87.9	(86.0 - 89.8)	91.9	(90.3 - 93.5) *
Ate fruits and vegetables > or = to 5 times				
Female	18.7	(16.7 - 20.7)	13.2	(10.3 - 16.1) *
Male	21.4	(20.0 - 22.8)	17.7	(14.9 - 20.5)
Total	20.1	(18.7 - 21.5)	15.5	(13.7 - 17.3) *
Drank > or = to 3 glasses of milk				
Female	11.6	(9.8 - 13.4)	10.6	(8.6 - 12.6)
Male	20.8	(18.5 - 23.1)	21.5	(18.9 - 24.1)
Total	16.2	(14.5 - 17.9)	16.2	(14.4 - 18.0)
Participated in currently recommended levels of physical activity				
Female	27.8	(25.5 - 30.1)	27.1	(23.5 - 30.7)
Male	43.8	(41.7 - 45.9)	37.2	(33.3 - 41.1) *
Total	35.8	(33.9 - 37.7)	32.2	(29.1 - 35.3)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Participated in previously recommended levels of physical activity				
Female	61.5	(59.2 - 63.8)	60.8	(57.0 - 64.6)
Male	75.8	(74.0 - 77.6)	70.9	(66.0 - 75.8)
Total	68.7	(67.1 - 70.3)	65.9	(62.5 - 69.3)
No vigorous physical activity				
Female	11.3	(10.0 - 12.6)	12.1	(10.2 - 14.0)
Male	7.9	(7.0 - 8.8)	9.0	(6.6 - 11.4)
Total	9.6	(8.7 - 10.5)	10.5	(8.9 - 12.1)
Enrolled in PE class				
Female	48.3	(42.9 - 53.7)	32.7	(25.5 - 39.9) *
Male	60.0	(55.7 - 64.3)	44.5	(36.7 - 52.3) *
Total	54.2	(49.4 - 59.0)	38.7	(31.5 - 45.9) *
Attended PE class daily				
Female	29.0	(23.4 - 34.6)	25.1	(17.9 - 32.3)
Male	37.1	(31.9 - 42.3)	31.2	(24.0 - 38.4)
Total	33.0	(27.7 - 38.3)	28.2	(21.3 - 35.1)
Exercised/played sports > 20 minutes during PE class				
Female	80.3	(76.2 - 84.4)	84.9	(78.1 - 91.7)
Male	87.2	(84.8 - 89.6)	87.7	(83.7 - 91.7)
Total	84.0	(81.0 - 87.0)	86.5	(82.5 - 90.5)
Played on > or = to 1 sports teams				
Female	50.2	(47.6 - 52.8)	54.7	(50.9 - 58.5)
Male	61.8	(59.1 - 64.5)	59.6	(56.2 - 63.0)
Total	56.0	(53.7 - 58.3)	57.1	(53.9 - 60.3)
Watched > or = to 3 hours per day of TV				
Female	36.3	(34.1 - 38.5)	29.6	(24.9 - 34.3)
Male	38.0	(35.5 - 40.5)	34.2	(30.2 - 38.2)
Total	37.2	(35.1 - 39.3)	31.9	(28.2 - 35.6)
At risk for becoming overweight				
Female	15.5	(14.1 - 16.9)	14.9	(12.9 - 16.9)
Male	15.8	(14.6 - 17.0)	13.6	(11.6 - 15.6)
Total	15.7	(14.8 - 16.6)	14.3	(13.0 - 15.6)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Overweight				
Female	10.0	(8.9 - 11.1)	9.2	(6.9 - 11.5)
Male	16.0	(14.9 - 17.1)	20.5	(17.0 - 24.0)
Total	13.1	(12.2 - 14.0)	15.0	(12.6 - 17.4)
Described themselves as overweight				
Female	38.1	(36.4 - 39.8)	37.2	(33.4 - 41.0)
Male	25.1	(23.9 - 26.3)	26.9	(23.4 - 30.4)
Total	31.5	(30.3 - 32.7)	31.9	(28.8 - 35.0)
Were trying to lose weight				
Female	61.7	(60.5 - 62.9)	63.1	(59.8 - 66.4)
Male	29.9	(28.3 - 31.5)	30.5	(27.5 - 33.5)
Total	45.6	(44.4 - 46.8)	46.5	(43.5 - 49.5)
Ate less food, fewer calories, or foods low in fat to lose weight or to keep from gaining weight				
Female	54.8	(53.2 - 56.4)	57.2	(53.1 - 61.3)
Male	26.8	(25.4 - 28.2)	27.0	(22.6 - 31.4)
Total	40.7	(39.5 - 41.9)	41.8	(38.1 - 45.5)
Went without eating for > or = to 24 hours to lose weight or to keep from gaining weight				
Female	17.0	(15.6 - 18.4)	17.8	(14.9 - 20.7)
Male	7.6	(6.7 - 8.5)	7.1	(5.3 - 8.9)
Total	12.3	(11.4 - 13.2)	12.4	(10.6 - 14.2)
Exercised to lose weight or to keep from gaining weight				
Female	67.4	(65.6 - 69.2)	73.3	(70.2 - 76.4) *
Male	52.9	(51.1 - 54.7)	51.6	(48.1 - 55.1)
Total	60.0	(58.6 - 61.4)	62.3	(59.4 - 65.2)
Took diet pills, powder, or liquids to lose weight or to keep from gaining weight				
Female	8.1	(6.6 - 9.6)	9.4	(7.0 - 11.8)
Male	4.6	(3.9 - 5.3)	4.2	(2.7 - 5.7)
Total	6.3	(5.3 - 7.3)	6.8	(5.3 - 8.3)
Vomited or took laxatives to lose weight or to keep from gaining weight				
Female	6.2	(5.4 - 7.0)	7.3	(5.3 - 9.3)
Male	2.8	(2.1 - 3.5)	3.7	(2.2 - 5.2)
Total	4.5	(4.0 - 5.0)	5.5	(4.2 - 6.8)

Table 1x. Percentage of high school students who engaged in health-risk behaviors, Youth Risk Behavior Survey, 2005 - United States and Indiana, Cont.

Health-risk behavior	National		Indiana	
	Percent	(95% CI)	Percent	(95% CI)
Lifetime Asthma				
Female	17.0	(15.7 - 18.3)	22.3	(18.6 - 26.0) *
Male	17.3	(15.9 - 18.7)	22.0	(19.1 - 24.9) *
Total	17.1	(16.2 - 18.0)	22.2	(20.0 - 24.4) *
Current Asthma				
Female	14.7	(13.6 - 15.8)	18.9	(16.0 - 21.8) *
Male	14.3	(13.0 - 15.6)	17.5	(14.8 - 20.2)
Total	14.5	(13.7 - 15.3)	18.2	(16.2 - 20.2) *
Asthma Episode or Attack				
Female	45.7	(41.4 - 50.0)	54.2	(47.6 - 60.8)
Male	30.4	(26.3 - 34.5)	27.0	(19.3 - 34.7)
Total	37.9	(35.0 - 40.8)	40.9	(34.7 - 47.1)
Described Health as Fair or Poor				
Female	9.6	(8.6 - 10.6)	8.9	(6.8 - 11.0)
Male	7.1	(6.4 - 7.8)	8.6	(6.4 - 10.8)
Total	8.3	(7.6 - 9.0)	8.8	(7.4 - 10.2)

*Indiana data significantly different from the national data.